

Isoscalar dipole strength in $^{208}_{82}\text{Pb}_{126}$: the spurious mode and the strength in the continuum

I.Hamamoto^{a,b,c} and H.Sagawa^d

^a Department of Mathematical Physics

Lund Institute of Technology at University of Lund, Lund, Sweden.

^b The Niels Bohr Institute

Blegdamsvej 17, DK-2100, Copenhagen Ø, Denmark.

^c Radiation Laboratory, RIKEN

Wako-shi, Saitama 351-0198, Japan.

^d Center for Mathematical Sciences

University of Aizu, Ikki-machi, Aizu-Wakamatsu, Fukushima 965, Japan.

ABSTRACT : Isoscalar dipole (compression) mode is studied first using schematic harmonic-oscillator model and, then, the self-consistent Hartree-Fock (HF) and random phase approximation (RPA) solved in coordinate space. Taking ^{208}Pb and the SkM* interaction as a numerical example, the spurious component and the strength in the continuum are carefully examined using the sum rules. It is pointed out that in the continuum calculation one has to use an extremely fine radial mesh in HF and RPA in order to separate, with good accuracy, the spurious component from intrinsic excitations.

PACS numbers : 21.10.Re, 21.60.Jz, 23.20.Js, 27.80.+w

1 Introduction

Though the isoscalar (IS) dipole resonance (a compression mode) was theoretically studied already more than 20 years ago [1, 2, 3, 4] and various experimental efforts to pin down the IS giant dipole resonance (ISGDR) have been made [5, 6, 7, 8, 9, 10], the distribution of the IS dipole strength obtained from the analysis of experimental data is still under dispute. The difficulty in extracting the experimental dipole strength comes from the ambiguity in the parameters of the optical potentials [11], which are needed in the analysis of the relevant (α, α') scattering data. In order to simplify our discussion, as a numerical example in the present article we discuss only the doubly magic nucleus $^{208}_{82}\text{Pb}_{126}$. A current hot issue in both theory and experiments is whether or not a considerable amount of IS dipole strength is found in the energy region of $E_x=8-17$ MeV, which is much lower than the so-called ISGDR with a peak around $E_x=23$ MeV. Available calculations have a tendency to give 1-3 MeV higher peak-energy of ISGDR than that identified so far in experiments.

The calculation of IS dipole strength requires special care since the spurious translation mode has to be fully separated from intrinsic excitations. In the self-consistent random-phase-approximation (RPA) calculation the spurious mode should appear at zero energy with a finite contribution to the energy-weighted sum-rule (EWSR). The response function to the IS dipole operator

$$D = \sum_k (r_k^3 Y_{1\mu}(\theta_k)) \quad (1)$$

contains the strengths of both the spurious mode and intrinsic excitations. In actual numerical calculations of self-consistent RPA the spurious mode appears not at zero energy but at either a finite energy or an imaginary energy, due to numerical inaccuracy as well as some possible practical approximations. This may lead to a question of whether or not the calculated IS dipole strength obtained in the energy region of intrinsic excitations is free from the spurious component. The difficulty in the present problem lies in the fact that a very small amount of the spurious component remaining in the energy region of intrinsic excitations can in turn produce an appreciable amount of the strength for

the operator (1) due to the power of r^3 . If the transition density coming from spurious component remaining in the region is proportional to $\frac{d\rho_0}{dr}$ where ρ_0 expresses the ground-state density, the IS dipole strength which is free from the spurious component is obtained by using the operator

$$\bar{D} = \sum_k (r_k^3 - \eta r_k) Y_{10}(\theta_k) \quad (2)$$

where $\eta = \frac{5}{3} \langle r^2 \rangle$ [3, 4, 12], instead of using that in (1). However, if some spurious components remain at the energies of various unperturbed particle-hole (ph) excitations, the relevant transition densities are similar to those of respective original ph excitations rather than $\frac{d\rho_0}{dr}$. If so, using the operator in (2) may not help to take away all strength coming from the spurious component remaining in the region of intrinsic excitations.

In ref. [12] we studied both the IS and isovector (IV) dipole mode using the self-consistent RPA calculation with the SkM* interaction, in which both the IS and IV correlations in RPA are taken into account solving both the Hartree-Fock (HF) and RPA equations in coordinate space using the Green's function. Examining the IS dipole strength obtained by using the operator in (2), in the energy region of $7 < Ex < 17$ MeV we obtained an appreciable amount of IS dipole strength almost exactly at the energies of various unperturbed ph excitations. Since the residual interaction relevant to the IS dipole correlation is in general attractive, the strength increased by the RPA correlation is expected to be found at energies lower than the unperturbed ones. Therefore, using the procedure described in ref. [12] we tried to remove the strength at finite energies, which seemed to come from the remaining spurious component. In the continuum at a given energy the excitations of various modes with various quantum numbers may coexist. The separation of the strengths coming from modes with different quantum numbers is easy, however, we have found no established way of separating the strengths originating from different modes with the same quantum numbers. Since both the spurious mode and IS dipole resonance are isoscalar and have the same spin-parity, $I^\pi = 1^-$, our basic idea used

in ref. [12] was the following. Near the isolated resonance, $E \approx E_n$, the relation

$$Im \left(G_{RPA}(\vec{r}, \vec{r}', E) \right) \propto \rho_n^{tr}(\vec{r}) \rho_n^{tr}(\vec{r}') \quad (3)$$

is known. At a high energy in the continuum, where resonances i and j with respective finite widths may overlap, we may write

$$\begin{aligned} Im \left(G_{RPA}(\vec{r}, \vec{r}', E) \right) &\propto a_i \rho_i^{tr}(\vec{r}) \rho_i^{tr}(\vec{r}') + b_j \delta \rho_j^{tr}(\vec{r}) \delta \rho_j^{tr}(\vec{r}') \\ &\equiv F(\vec{r}, \vec{r}') \end{aligned} \quad (4)$$

where the modes, i and j , have the same spin-parity and are diagonal in RPA. Choosing some $f(\vec{r})$ so that

$$\int d\vec{r}' \rho_i^{tr}(\vec{r}') f(\vec{r}') = 0 \quad (5)$$

one obtains

$$\delta \rho_j^{tr}(\vec{r}) \propto \int d\vec{r}' F(\vec{r}, \vec{r}') f(\vec{r}') \quad (6)$$

Choosing $f(\vec{r}) = r Y_{1\mu}$ in the IS dipole case, we obtained $\delta \rho_j^{tr}(\vec{r})$, which was supposed to be the transition density at the energy E , that carried the spurious $r Y_{1\mu}$ strength. Then, we subtracted $\delta \rho_j^{tr}(\vec{r})$ from the transition density $\rho_{tr}(\vec{r})$ at E for the operator D in (1),

$$\rho_{tr}(\vec{r}) \propto \int d\vec{r}' Im \left(G_{RPA}(\vec{r}, \vec{r}', E) \right) (r')^3 Y_{1\mu}(\theta'_k) \quad (7)$$

Using the above subtraction procedure, it seemed that in ref. [12] we subtracted a bit too much strength as was criticized, for example, in ref. [13]. The oversubtraction happened, since the remaining ph excitations which carry some spurious strength were not really RPA solutions.

In all available calculations of ISGDR except those in refs. [3, 12, 18] the states in the continuum are approximated using the expansion in terms of discrete basis. Namely, the continuum wave functions are expanded in terms of harmonic oscillator basis with a maximum principal-quantum-number N_{max} . In those discrete-basis calculations the operator in (2) is commonly used in order to subtract the spurious components. The IS

dipole strength obtained in the energy region of $E_x = 8 - 17$ MeV depends very much on publications. The relativistic RPA calculation in ref. [17] produced especially a large amount of the IS dipole strength in the low-energy region, for all effective interactions used. In none of the discrete-basis calculations the EWSR carried by the spurious state and intrinsic excitations is carefully examined.

In the present paper we use EWSR as a measure of the validity of numerical calculations. The EWSR for the IS dipole operator in (1) is written as

$$\begin{aligned} S(r^3 Y_{10}) &\equiv \sum_n E_n B(D; 0 \rightarrow 1^- n) \\ &= \left(\frac{3}{4\pi}\right) \left(\frac{\hbar^2 A}{2M}\right) 11 \langle r^4 \rangle \end{aligned} \quad (8)$$

where E_n expresses the energy of RPA states, while $\langle r^4 \rangle$ on the r.h.s. is evaluated with the HF ground state [14]. The total EWSR in (8) consists of the contribution by the spurious state S_{spr} and that by the intrinsic excitations S_{intr} , $S = S_{spr} + S_{intr}$. The latter can be written as [4]

$$\begin{aligned} S_{intr}(r^3 Y_{10}) &= \sum_{intrinsic, i} E_i B(D; 0 \rightarrow 1^- i) \\ &= \left(\frac{3}{4\pi}\right) \left(\frac{\hbar^2 A}{2M}\right) \left(11 \langle r^4 \rangle - \frac{25}{3} \langle r^2 \rangle^2\right) \end{aligned} \quad (9)$$

which is indeed equal to the EWSR for the IS dipole operator in (2).

In sect. 2 we use a harmonic oscillator model to evaluate the distribution of the EWSR of IS dipole strength among the spurious state, the lower-lying and higher-lying intrinsic excitations. This schematic model is very much simplified, nevertheless, we expect that the semi-quantitative feature of the result obtained from the model should remain unchanged in realistic calculations. In sect. 3 we present the result of the self-consistent RPA calculation with the Skyrme SkM* interaction on ^{208}Pb , which is the continuum calculation solving both HF and RPA in coordinate space. The result is compared with discrete-basis calculations, in which the wave functions in the continuum are expanded in terms of harmonic oscillator basis. In sect.4 conclusion and discussions are given.

2 Harmonic oscillator model

The distribution of IS dipole strength may be studied in terms of a simplified model [15] in which the particle motion is described by a harmonic oscillator potential. The particle excitations produced by the field (1) are governed by the selection rule $\Delta N=1$ or 3 and have the energies $\hbar\omega_0$ and $3\hbar\omega_0$, respectively. For a single completed shell with principal quantum number N , the transition strength is given by

$$\sum_{\nu_N, \nu_{N+1}} |\langle \nu_{N+1} | r^3 Y_{10} | \nu_N \rangle|^2 = \frac{3}{4\pi} \left(\frac{\hbar}{2M\omega_0} \right)^3 \frac{1}{12} (N+1)(N+2)(N+3)(13N^2 + 52N + 50) \quad (10)$$

$$\sum_{\nu_N, \nu_{N+3}} |\langle \nu_{N+3} | r^3 Y_{10} | \nu_N \rangle|^2 = \frac{3}{4\pi} \left(\frac{\hbar}{2M\omega_0} \right)^3 \frac{1}{12} (N+1)(N+2)(N+3)(N+4)(N+5) \quad (11)$$

where ν_N represents the quantum numbers needed to specify the single-particle states in the shell N . For simplicity, we do not include the spin-isospin degeneracy in the present model. Since the last three occupied shells contribute to the total $\Delta N=3$ strength we obtain

$$\sum_{N=N_F-2}^{N_F} \sum_{\nu_N, \nu_{N+3}} |\langle \nu_{N+3} | r^3 Y_{10} | \nu_N \rangle|^2 = \frac{3}{4\pi} \left(\frac{\hbar}{2M\omega_0} \right)^3 \frac{1}{12} (N_F+1)(N_F+2)(N_F+3)(3N_F^2 + 12N_F + 20) \quad (12)$$

while only the last-filled $N = N_F$ shell contributes to the $\Delta N=1$ strength.

Now, the major part of the $\Delta N=1$ strength, which is obtained by setting $N = N_F$ in (10), corresponds to the spurious excitation, namely center of mass motion, while in the present harmonic oscillator model the $\Delta N=3$ strength contains no spurious component. Subtracting the strength coming from the spurious excitation

$$\frac{3}{4\pi} \left(\frac{\hbar}{2M\omega_0} \right)^3 \frac{25}{24} (N_F + 1)(N_F + 2)^3 (N_F + 3), \quad (13)$$

from the expression (10) with $N = N_F$, we obtain the strength of the intrinsic $\Delta N=1$ excitation

$$\frac{3}{4\pi} \left(\frac{\hbar}{2M\omega_0} \right)^3 \frac{1}{24} N_F (N_F + 1)(N_F + 2)(N_F + 3)(N_F + 4) \quad (14)$$

We note that the total number of particles in the present model is written as

$$\frac{1}{6}(N_F + 1)(N_F + 2)(N_F + 3) \quad (15)$$

while

$$\sum_{N=0}^{N_F} \langle r^2 \rangle = \left(\frac{\hbar}{2M\omega_0} \right) \frac{1}{4}(N_F + 1)(N_F + 2)^2(N_F + 3) \quad (16)$$

and

$$\sum_{N=0}^{N_F} \langle r^4 \rangle = \left(\frac{\hbar}{2M\omega_0} \right)^2 \frac{1}{2}(N_F + 1)(N_F + 2)(N_F + 3)(N_F^2 + 4N_F + 5) \quad (17)$$

The spurious contribution (13) is estimated by recognizing that the spurious component is found only at the $\Delta N=1$ excitations and that the transition density of the spurious excitation is proportional to the radial derivative of the ground-state density, $d\rho_0/dr$.

From the expressions, (12) and (14), one obtains the ratio of the unperturbed transition strength of $\Delta N=1$ excitations to that of $\Delta N=3$ ones to be 1:6 for $N_F \gg 1$. Correspondingly, the ratio of the unperturbed EWSR is 1:18.

Introducing the residual interaction of a separable form

$$V = c D^\dagger D \quad (18)$$

where c expresses an attractive coupling constant, we solve the RPA equation and estimate the distribution of the transition strength together with the RPA energies. The following results of cases (a) and (b) are summarized in Table I.

(a) If we choose c so that the higher-lying RPA solution is obtained at $(2.5)\hbar\omega_0$, the lower-lying RPA solution is found at $(0.872)\hbar\omega_0$. Then, the RPA transition strength is distributed with 1.09:2.79 over the lower-lying and higher-lying RPA solutions, while the distribution of the EWSR is 0.95:6.98 .

(b) If we choose c so that the higher-lying RPA solution is obtained at $(2.8)\hbar\omega_0$, the lower-lying RPA solution is found at $(0.961)\hbar\omega_0$. Then, the RPA transition strength is distributed with 0.60:2.62 between the lower-lying and higher-lying RPA solutions, while the distribution of the EWSR is 0.58:7.34 .

Comparing the peak energies of the RPA solutions with those obtained by available self-consistent HF plus RPA calculations, one expects that the realistic situation may lie somewhere between the case (a) and (b). Before the spurious component $\delta \rho_j^{tr}(\vec{r})$ in (6) was subtracted in our self-consistent RPA calculations of ref. [12], a number of lower-energy peaks at $E_x \lesssim 17$ MeV for the operator (2) were found nearly at respective unperturbed ph energies. In contrast, the higher-energy peak around 25 MeV was collectively constructed by shifting the unperturbed strength coherently from the region of higher energy. In other available self-consistent RPA calculations, for example in refs. [16] and [17] in which the expansion in terms of harmonic oscillator basis was used, or in ref. [18] which was a continuum calculation of relativistic Hartree plus RPA calculation, a considerable amount of low-energy strength appeared also around unperturbed ph energies. Compared with those HF plus RPA calculations, in the case (a) of Table I the higher-lying RPA peak is pushed down by about the same order of magnitude and the lower-lying peak is too strongly pushed down, while in (b) the higher-lying peak is appreciably less pushed down and the lower-lying peak is still slightly lower.

On the other hand, one might wonder whether or not the ratio of the coupling between the $\Delta N=1$ and 3 excitations to that within the excitations of a given ΔN is realistic in the present schematic interaction of (18). In the present harmonic-oscillator model the effective strength of the coupling between the $\Delta N=1$ and 3 excitations can be measured from the increase of EWSR in the lower-lying RPA solution compared with that in the unperturbed $\Delta N=1$ excitation. The portion of the EWSR in (8) carried by the lower-energy peak is increased from 2.3 to 5.2 percent in the case (a) and to 3.2 percent in the case (b), as shown in Table I.

3 Hartree-Fock and RPA calculations with Skyrme interactions

In the HF calculation with the SkM* interaction for ^{208}Pb we obtain

$$\begin{aligned}\langle r^2 \rangle &= 30.85 \text{ fm}^2 \\ \langle r^4 \rangle &= 1218.2 \text{ fm}^4\end{aligned}\tag{19}$$

Thus, using eqs. (8) and (9) the EWSR becomes

$$S(r^3 Y_{10}) = 13.79 \times 10^6 \text{ fm}^6 \text{ MeV}\tag{20}$$

$$S_{intr}(r^3 Y_{10}) = 5.632 \times 10^6 \text{ fm}^6 \text{ MeV}\tag{21}$$

From eqs. (20) and (21) the contribution by the spurious state to the EWSR for the operator $r^3 Y_{1\mu}$ is $8.16 \times 10^6 \text{ fm}^6 \text{ MeV}$, which is 59.2 percent of the total EWSR in (20).

In the present work we perform the self-consistent RPA calculation with the SkM* interaction, in which both the IS and IV correlations in RPA are taken into account solving both HF and RPA equations in coordinate space using the Green's function. The difference of the present calculation from that in ref. [12] is that we use the radial mesh (Δr) of 0.1 fm in both the HF and RPA calculations and, moreover, we carefully check the sum-rules consumed by both intrinsic and spurious excitations. We present the numerical result without adopting the subtraction procedure, which was used in ref. [12] and was also explained in the Introduction. In ref. [12] $\Delta r=0.1 \text{ fm}$ was used in HF, while we adopted $\Delta r=0.3 \text{ fm}$ in RPA. In Fig. 1 we show the RPA response functions, which are smeared out using the width of 0.5 MeV , for the operator (1) obtained with $\Delta r=0.1 \text{ fm}$ and 0.3 fm and for the operator (2) calculated with $\Delta r=0.1 \text{ fm}$. A great, somewhat unexpected improvement in the result of the present calculation with a finer radial mesh in RPA is : the appreciable difference between the response functions to the operators (1) and (2) in the energy region of the higher peak around 25 MeV is drastically decreased. Moreover, the spurious state appears now slightly below 0 MeV (namely, at an imaginary energy) and consumes a much larger portion of EWSR, S_{spr} .

As in ref. [12], our effective interaction used in RPA consists only of the part in the spin-independent channels. Using discrete-basis calculations, in which the inclusion of the full Skyrme effective interaction is straightforward and easy, we have checked that the RPA response function of the present IS dipole operator changes very little when we use the full effective interaction, as seen in Tables II and III.

In Table II we show the transition strength for the operator (1), the contribution to EWSR, and the percent of the total EWSR, $S(r^3Y_{10})$, at the spurious state and in the region of $2 < E_x < 17$ MeV, $17 < E_x < 70$ MeV, and $E_x > 70$ MeV. In Table III the same quantities are tabulated for the operator (2). The borderline 17 MeV is chosen, since almost all $\Delta N = 1, I^\pi = 1^-$ ph excitations in the HF potential lie below 17 MeV and no unperturbed 1^- ph excitations are found for $18 < E_x < 24$ MeV.

As is previously known, it is seen from Table II that the portion of the EWSR found in the energy region of intrinsic excitations ($E_x > 2$ MeV) clearly shows the presence of the spurious components remaining in the region. From Table III it is seen that the ratio of the EWSR lying in the region of $E_x < 17$ MeV to that for $E_x > 17$ MeV is 1:4.9. More importantly, the portion of the EWSR obtained for $E_x > 2$ MeV is 39.1 percent of the total EWSR, compared with the full value, 40.8 percent. It should be mentioned that in the continuum calculation it is technically very difficult to obtain accurate values of the transition strength carried by very sharp peaks including the spurious state, in contrast to the discrete-basis calculations. Furthermore, in the present continuum calculation the spurious state appears slightly below 0 MeV. Thus, reducing the effective ph interactions by multiplying a factor of 0.9902, we obtain the spurious state at $E_x=0.107$ MeV, from which we estimate the carried transition strength. Though it is clear that the continuum calculations performed using a much finer radial mesh than 0.1 fm would further improve the accuracy, here we do not pursue it any further.

In Fig. 2 we show the calculated RPA response functions to the operators, (1) and (2), as a function of excitation energy. Though the peaks at $E_x > 7.5$ MeV lie in the

continuum, calculated lower-lying peaks are too sharp to be plotted. Since the spreading width is anyhow not included in the RPA response function, in Fig. 2 we plot the quantities, which are obtained by smearing out the calculated RPA response function with the width of 1 MeV. It is seen that an appreciable amount of IS dipole strength is present in the energy region of $E_x=7-17$ MeV, which is much lower than the so-called ISGDR around 25 MeV. The presence of the lower-lying IS dipole strength indicates that the ISGDR appearing around 25 MeV is not really a good collective mode in the nuclear system.

For reference, in Tables II and III we show also the result of discrete-basis calculations for the operators, (1) and (2). How to truncate the space of harmonic oscillator basis seems to depend on publications. What we have adopted in the present work, which is along the line of ref. [16], is the following. First we construct the HF potential, which is calculated in coordinate space and, then, calculate the one-particle energies and wave-functions for the potential using the harmonic oscillator basis with the radial node $n^{HO} \leq n_{max}^{HO}$, where n_{max}^{HO} is taken to be independent of ℓ . Then, the RPA equation is solved including the lowest-lying $\frac{1}{2}(N_{max}^{HF} - \ell)$ one-particle levels for a given ℓ , where N_{max}^{HF} is taken to be independent of ℓ . This truncation of the ph space included in RPA is approximately the same as that limiting the ph space by the ℓ -independent maximum energy of particle configurations. It is noted that, as n_{max}^{HO} increases, the bound one-particle energies and wave-functions approach the eigen energies and eigen functions of the HF potential, respectively, while the properties of the positive-energy one-particle levels can be far away from the correct ones unless n_{max}^{HO} is taken to be ∞ . In actual calculations one must take finite values of n_{max}^{HO} and N_{max}^{HF} and, thus, it is not guaranteed that using finite larger values of n_{max}^{HO} and N_{max}^{HF} produces a more accurate result of IS dipole strength.

For the parameter set, $n_{max}^{HO} = 12$ and $N_{max}^{HF} = 24$, in Tables II and III we show both the result obtained by using the full effective interaction in RPA and the one by employing the same effective interaction used in our continuum RPA calculation. The fact that very

little difference is found between these two results indicates that the IS dipole strength obtained from our continuum RPA calculation is reliable. Comparing the numbers in the last column of Table II with those in Table III, in the discrete-basis calculation the difference between the strengths calculated with the operators (1) and (2) is relatively minor in the region of $E_x > 17$ MeV. On the other hand, from Table II we notice that the calculated energy of the spurious state is pretty high compared with the continuum calculation and, furthermore, only 92 percent of the correct portion of EWSR is consumed by the calculated spurious state. The properties of the spurious state converge surprisingly slowly to the right ones, as we increase n_{max}^{HO} and N_{max}^{HF} . On the other hand, if we use a value of n_{max}^{HO} larger than 12, the calculated response function in the continuum starts to behave strangely instead of converging to some reasonable result. This strange behavior of discrete-basis calculations seems to be known for experts in the fields [19].

4 Conclusion and discussions

First, the distribution of IS dipole strength for heavy nuclei is studied using the schematic harmonic-oscillator model, in which the strength is distributed among the spurious mode, the low-energy and high-energy peaks. Though 96 percent of the unperturbed $\Delta N = 1$ strength for the operator $r^3 Y_{1\mu}$ corresponds to the spurious excitation, the remaining 4 percent contributes to the intrinsic excitations. The spurious state consumes 56.8 percent of the total EWSR for the operator $r^3 Y_{1\mu}$, while the low-energy RPA peak may carry up till 5 percent of the EWSR.

Secondly, the result of IS dipole strength in ^{208}Pb is presented, which is calculated using the self-consistent HF plus RPA solved in coordinate space using the Green's function method, with better accuracy than in ref. [12]. If one performs numerical calculations with very good accuracy, the IS dipole strength of intrinsic excitations should not depend on whether the operator (1) or (2) is used. At the same time, the spurious mode should appear close to zero energy and consume the whole portion of EWSR expected. In our

present continuum calculation we could not attain such a satisfactory accuracy. On the other hand, if the calculated strengths with the operators (1) and (2) are appreciably different from each other in the energy region of intrinsic excitations, the spurious component remaining in the region may not have the transition density proportional to $\frac{d\rho_0(r)}{dr}$. In the present work we have not attempted to take away this kind of possible spurious component from the region of intrinsic excitations, since we find no reliable way of excluding it.

Using the radial mesh of 0.1 fm in both the HF and RPA calculations, the spurious state is obtained slightly below 0 MeV. Multiplying the effective ph interaction by a factor of 0.9902, we push up the energy to 0.107 MeV and estimate the EWSR carried by the state, which turns out to be the major portion of the expected EWSR. On the other hand, the EWSR carried by the calculated states for $E_x > 2$ MeV is 114 and 96 percent of the expected EWSR for the operators (1) and (2), respectively. The accuracy of the continuum calculations is appreciably improved by decreasing Δr used in solving RPA from 0.3 fm in ref. [12] to 0.1 fm , however, it is still away from being very satisfactory. In the continuum calculations the accuracy of separating the spurious component from intrinsic excitations is surprisingly sensitive to the size of radial mesh employed in RPA. Thus, the response function of IS dipole strength has to be calculated using a very fine radial mesh, in order to obtain a reliable numerical result. From the present study we find that slightly more than 5 percent of the total EWSR is expected as intrinsic excitations lying in the low-energy region below $E_x < 17$ MeV.

The expansion of HF and RPA solutions in terms of harmonic oscillator basis can work for bound states if one includes a sufficiently large number of bases [20]. Thus, in the present study of IS dipole strength the properties of the spurious mode become better and better as the number of bases increases, though we find that the convergence of the properties is unexpectedly slow as n_{max}^{HO} and N_{max}^{HF} increase. On the portion of EWSR found in the continuum the discrete-basis calculation presented in Table II and

Fig. 1 produces 9 percent larger in the $2 < E_x < 17$ MeV region and 6 percent smaller for $E_x > 17$ MeV, compared with our continuum calculation. However, the real problem is : If we further enlarge the space of discrete-basis, the calculation starts to produce more than 100 percent of EWSR, which are certainly unreasonable. Thus, it is not clear whether or not the discrete-basis calculations can be reliably used for the estimate of IS dipole strength.

The authors are grateful to Drs. N. Van Giai and G. Colo for the generous advice on the discrete-basis calculation. I.H. acknowledges the financial supports provided by Crafoordska Stiftelsen and RIKEN. H.S. acknowledges financial support by the Japanese Ministry of Education, Science, Sports and Culture by Grant-In-Aid for Scientific Research under the program number C(2) 12640284.

References

- [1] T. J. Deal, Nucl. Phys. **A217**, 210 (1973).
- [2] M. N. Harakeh, Phys. Lett. **90B**, 13 (1980); M. N. Harakeh and A. E. L. Dieperink, Phys. Rev. **C23**, 2329 (1981).
- [3] N. Van Giai and H. Sagawa, Nucl. Phys. **A371**, 1 (1981).
- [4] T. S. Dumitrescu and F. E. Serr, Phys. Rev. **C27**, 811 (1983).
- [5] H. P. Morsch, M. Rogge, P. Turek, and C. Mayerboricke, Phys. Rev. Lett. **45**, 337 (1980).
- [6] C. Djalali, N. Marty, M. Morlet, and A. Willis, Nucl. Phys. **A380**, 1 (1982).
- [7] G. S. Adams, T. A. Carey, J. B. McClelland, J. M. Moss, S. J. Seestrom-Morris, and D. Cook, Phys. Rev. **C33**, 2054 (1986).

- [8] B. F. Davis *et al.*, Phys. Rev. Lett. **79**, 609 (1997).
- [9] H. L. Clark, Y.-W. Lui, D. H. Youngblood, K. Bachtr, U. Garg, M. N. Harakeh, and N. Kalantar-Nayestanaki, Nucl. Phys. **A649**, 57c (1999).
- [10] H. L. Clark, Y.-W. Lui, and D. H. Youngblood, Phys. Rev. **C63**, 031301(R) (2001).
- [11] H. L. Clark, Y.-W. Lui, and D. H. Youngblood, Nucl. Phys. **A687**, 80 (2001).
- [12] I. Hamamoto, H. Sagawa, and X. Z. Zhang, Phys. Rev. **C57**, R1064 (1998).
- [13] S. Shlomo and A. I. Sanzhur, Phys. Rev. **C65**, 044310 (2002).
- [14] D. J. Thouless, Nucl. Phys. **22**, 78 (1961).
- [15] A. Bohr and B. R. Mottelson, *Nuclear Structure* (Benjamin, Reading, MA, 1975), Vol. II, Chapter 6.
- [16] G. Colo, N. Van Giai, P. F. Bortignon, and M. R. Quaglia, Phys. Lett. **B485**, 362 (2000).
- [17] D. Vretenar, A. Wandelt, and P. Ring, Phys. Lett. **B487**, 334 (2000).
- [18] J. Piekarewicz, Phys. Rev. **C62**, 051304 (2000).
- [19] N. Van Giai and G. Colo, private communications.
- [20] J. P. Blaizot and D. Gogny, Nucl. Phys. **A284**, 429 (1977).

Figure captions

Figure 1 : RPA response functions to the IS dipole operators, (1) and (2), as a function of excitation energy, which are obtained from the self-consistent HF plus RPA calculations solved in coordinate space. A radial mesh $\Delta r=0.1 \text{ fm}$ is used in HF, while the results obtained by using $\Delta r=0.1$ and 0.3 fm in RPA are compared. We show the calculated response functions, which are smeared out using the width of 0.5 MeV .

Figure 2 : RPA response functions to the IS dipole operators, (1) and (2), as a function of excitation energy, which are obtained from the self-consistent continuum calculation (CRPA) with $\Delta r=0.1 \text{ fm}$ and the discrete-basis calculation with $n_{max}^{HO} = 12$ and $N_{max}^{HF} = 24$ (DRPA). In all calculations the $(1, \tau \cdot \tau)$ effective interaction (see the caption to Table II) is used. In order to see more easily the total distribution of the strength, we plot both CRPA and DRPA response functions, which are smeared out using the width of 1 MeV .

Table 1: Energy, reduced transition strength $B(D)$, and contribution to the EWSR for the operator $r^3Y_{1\mu}$, which are calculated in the harmonic oscillator model for $N_F \gg 1$. B(D) and EWSR are given in arbitrary unit.

Case	energy in $\hbar\omega_0$	B(D)	EWSR (%)
unperturbed	spurious (1.0)	$\frac{25}{24} = 1.042$	$\frac{25}{24} = 1.042$ (56.8)
	1.0	$\frac{1}{24} = 0.042$	$\frac{1}{24} = 0.042$ (2.3)
	3.0	$\frac{1}{4} = 0.25$	$\frac{3}{4} = 0.75$ (40.9)
(a)	0.0	∞	1.042 (56.8)
	0.872	0.109	0.095 (5.2)
	2.5	0.279	0.698 (38.0)
(b)	0.0	∞	1.042 (56.8)
	0.961	0.060	0.058 (3.2)
	2.8	0.262	0.734 (40.0)

Table 2: Energy, reduced transition strength $B(D)$, and contribution to the EWSR for the operator $r^3Y_{1\mu}$ in ^{208}Pb , which are calculated in the self-consistent RPA using the SkM* interaction. Inside the bracket of the forth column the ratio to the total EWSR for the operator $r^3Y_{1\mu}$, (20), is expressed in percent. For discrete calculations with $n_{max}^{HO} = 12$ and $N_{max}^{HF} = 24$ we show two sets of calculated results, the one obtained by using the full effective interaction in RPA and the other calculated by using only the spin-independent part as that employed in our continuum RPA calculation. The former is denoted by “full”, while the latter is marked by “(1, $\tau \cdot \tau$)”. In the continuum calculation it is technically very difficult to estimate an accurate value of the transition strength carried by the spurious state, which is given with “ $\approx$ ” in the third and fourth column. See the text for details.

Type of calculation	energy in MeV	B(D) in fm^6	EWSR in fm^6MeV (%)
correct calculation	0.0	∞	8.16×10^6 (59.2)
	$0 < E_x$		5.63×10^6 (40.8)
continuum	0.107	$\approx 7.5 \times 10^7$	$\approx 8.0 \times 10^6$
(1, $\tau \cdot \tau$)	$2 < E_x < 17$	1.275×10^5	1.433×10^6 (10.4)
	$17 < E_x < 70$	1.973×10^5	4.856×10^6 (35.2)
	$70 < E_x < 150$	1.3×10^3	1.14×10^5 (0.83)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 16$)	1.401	5.342×10^6	7.485×10^6 (54.7)
full	$2 < E_x < 17$	1.347×10^5	1.534×10^6 (11.1)
	$17 < E_x < 70$	1.911×10^5	4.677×10^6 (33.9)
	$70 < E_x < 88$	52.8	3.869×10^3 (0.028)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 24$)	0.758	9.949×10^6	7.541×10^6 (54.7)
full	$2 < E_x < 17$	1.346×10^5	1.527×10^6 (11.1)
	$17 < E_x < 70$	1.914×10^5	4.683×10^6 (34.0)
	$70 < E_x < 190$	476.8	4.141×10^4 (0.30)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 24$)	0.406	1.850×10^7	7.538×10^6 (54.7)
(1, $\tau \cdot \tau$)	$2 < E_x < 17$	1.411×10^5	1.560×10^6 (11.3)
	$17 < E_x < 70$	1.899×10^5	4.653×10^6 (33.7)
	$70 < E_x < 190$	475.6	4.140×10^4 (0.30)

Table 3: The same quantities as in Table II, but for the operator $(r^3 - \eta r)Y_{1\mu}$ in ^{208}Pb . See the caption to Table II.

Type of calculation	energy in MeV	$B(\bar{D})$ in fm^6	EWSR in $fm^6 MeV$ (%)
correct calculation	0.0 $0 < E_x$	0.0	0.0 (0) 5.63×10^6 (40.8)
continuum (1, $\tau \cdot \tau$)	0.107 $2 < E_x < 17$ $17 < E_x < 70$ $70 < E_x < 150$	1.163×10^4 8.294×10^4 1.802×10^5 0.005×10^5	1.240×10^3 (0.009) 9.064×10^5 (6.6) 4.440×10^6 (32.2) 0.042×10^6 (0.3)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 16$) full	1.401 $2 < E_x < 17$ $17 < E_x < 70$ $70 < E_x < 88$	1.672×10^3 9.369×10^4 1.878×10^5 36.2	2.342×10^3 (0.011) 1.006×10^6 (7.3) 4.580×10^6 (33.2) 2.653×10^3 (0.019)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 24$) full	0.758 $2 < E_x < 17$ $17 < E_x < 70$ $70 < E_x < 190$	2.005×10^3 9.492×10^4 1.889×10^5 368.1	1.520×10^3 (0.011) 1.015×10^6 (7.4) 4.604×10^6 (33.4) 3.165×10^4 (0.23)
discrete ($n_{max}^{HO} = 12, N_{max}^{HF} = 24$) (1, $\tau \cdot \tau$)	0.406 $2 < E_x < 17$ $17 < E_x < 70$ $70 < E_x < 190$	3.607×10^3 9.544×10^4 1.886×10^5 364.7	1.464×10^3 (0.011) 1.023×10^6 (7.4) 4.597×10^6 (33.3) 3.130×10^4 (0.23)



